

Energy Conditions and Segre Classification of Phantom Fields

Janilo Santos*

Universidade Federal do Rio Grande do Norte, Departamento de Física, C.P. 1641, 59072-970 Natal, RN, Brasil

J. S. Alcaniz†

Observatório Nacional, Rua Gal. José Cristino 77, 20921-400 Rio de Janeiro - RJ, Brasil

(Dated: February 2, 2008)

Recent discoveries in the field of observational cosmology have provided increasing evidence that the Universe is undergoing a late time acceleration, which has also stimulated speculations on the nature of the dark component responsible for such a phenomenon. Among several candidates discussed in the current literature, phantom fields, an exotic scalar field with a negative kinetic term and that violates most of the classical energy conditions, appear as a real possibility according to recent observational analysis. In this paper we examine the invariant characterization for the energy-momentum tensor of phantom fields through the Segre algebraic classification in the framework of general relativity. We also discuss some constraints which are imposed on the values of $V(\phi)$ from the classical energy conditions.

PACS numbers: 98.80.Jk; 98.80.-k; 04.20.Cv

I. INTRODUCTION

The algebraic classification of symmetric second-order tensors locally defined on a 4-dimensional Lorentzian manifold, such as the Ricci tensor R_{ab} , the Einstein tensor G_{ab} and the energy-momentum tensor T_{ab} , is known as Segre classification [1]. The major idea underlying most of classifications in science is the concept of equivalence. Clearly the objects may be classified in different ways according to the criteria one chooses to group them into equivalence classes. There are, however, criteria that prove to be more important than others. The great appeal of Segre classification in general relativity is that it incorporates, ab initio, the Lorentzian character of space-time. It is of interest in several contexts such as, for example, in understanding purely geometrical features of space-times [2], in classifying and interpreting matter fields in general relativity [3] and in higher dimensional theories [4] (e.g., 5-D brane-worlds [5]) or still as part of the procedure for checking whether apparently different space-times are in fact locally the same up to coordinate transformations (equivalence problem [6]). Because of Einstein's equations $G_{ab} = \kappa T_{ab}$ the Einstein tensor (and so R_{ab}) has the same algebraic classification as the energy-momentum tensor. This can be used to decide, on base of generic features such as the Segre class, which energy-momentum tensors do couple to a given geometry.

A great deal of difficulty in Segre classifying second order tensors in the context of general relativity is that while G_{ab} and R_{ab} are universal functions of the space-time geometry, T_{ab} depends on the symmetries of the model as well as on the particular type or combination of matter fields present in the cosmological scenario. In this

regard, a powerful way of imposing physical constraints and searching realistic forms for the energy-momentum tensor is through cosmological observations. The high degree of isotropy observed in the Cosmic Microwave Background Radiation [7], for instance, restricts the general form of T_{ab} while results from distance measurements using type Ia supernovae (which constitute the most direct evidence for the current cosmic acceleration) [8] impose important constraints on the physical quantities of the cosmic fluid.

In light of an impressive convergence of observational results, several groups have recently tested the viability of different matter fields (or, equivalently, different forms of T_{ab}) as a realistic description for the matter content of our Universe. The two favorite candidates are the energy density stored on the true vacuum state of all existing fields in the Universe, i.e., the cosmological constant ($T_{ab}^{\Lambda} = \Lambda g_{ab}$) and the potential energy density $V(\phi)$ associated with a dynamical scalar field ϕ , usually called dynamical dark energy (see, e.g., [9] for a recent review on this topic). Physically, the role of such a dynamical dark energy can be played by any scalar field violating the so-called strong energy condition (see Sec. III for a discussion). Two possibilities, however, have been exhaustively explored in the current literature, namely, quintessence scalar fields with positive kinetic term and an equation-of-state parameter $w \geq -1$ [10], and phantom fields with negative kinetic term and $w < -1$ [11, 12, 13]. In terms of the parameter w , the former case can be seen as a very general scenario, which includes cold dark matter models with a cosmological constant (Λ CDM), $w = -1$, and cosmologies dominated by topological defects (e.g., domain walls, strings and textures) for which $w = -\frac{n}{n-2}$, n being the dimension of the defect. The latter case in turn is characterized by strange properties such as, for instance, the fact that its energy density increases with the expansion of the Universe (in contrast with quintessence fields); the possibility of a rip-off of large and small scale

*Electronic address: janilo@dfte.ufrn.br

†Electronic address: alcaniz@on.br

structures of matter; a possible occurrence of future curvature singularity, etc. [12]. Although having these unusual characteristics, a phantom type behavior is predicted by several scenarios, e.g., kinetically driven models [14] and some versions of brane world cosmologies [15]. From the observational point of view, phantom dark energy is found to be compatible with most of the classical cosmological tests and provide a better fit to type Ia supernovae observations than do Λ CDM or quintessence scenarios ($w > -1$). Therefore this means that, although exotic, phantom fields may be the dominant form of energy in our universe.

In this paper we examine the invariant characterization for the energy-momentum tensor of phantom fields through the Segre algebraic classification in the framework of general relativity. In Sec. II, by using real null tetrad technique, we show that phantom fields can be classified in two different subclasses of equivalence which are represented by Segre types $[1, (111)]$ and $[(1, 111)]$. In Sec. III we present the so-called energy conditions of general relativity for the whole Segre class $[1, 111]$ and also examine the further restrictions they impose on the values of the phantom field potential $V(\phi)$. We end this paper by summarizing the main results in the conclusion Section.

II. SEGRE CLASSIFICATION

A classification of a generic symmetric second order tensor T_{ab} can be cast in terms of the eigenvalue problem

$$(T^a_b - \lambda \delta^a_b) v^b = 0 \quad (1)$$

where λ are eigenvalues, v^a are eigenvectors, and the mixed tensor T^a_b may be thought of as a linear map $T_p(M) \mapsto T_p(M)$. M is a real 4-dimensional space-time manifold locally endowed with a Lorentzian metric of signature $(-+++)$, $T_p(M)$ denotes the tangent space to M at a point $p \in M$ and latin indices range from 0 to 3. Because of the Lorentzian character of the metric the mixed form of the energy-momentum tensor is no more symmetric and T^a_b may not have a diagonal matrix representation, i.e., it need not have four linearly independent eigenvectors. However, using the *Jordan canonical forms* of the matrix of an operator, and imposing the Lorentzian character of the metric on M , it has been shown [1, 3] (for detailed calculations in $n \geq 5$ dimensional space-times and a review on this topic see [20]) that any energy-momentum tensor defined on $T_p(M)$ re-

duces to one of the four canonical forms:

$$\begin{aligned} [1, 111] \quad T_{ab} &= 2\sigma_1 l_{(a} m_{b)} + \sigma_2 (l_a l_b + m_a m_b) \\ &\quad + \sigma_3 x_a x_b + \sigma_4 y_a y_b \\ [211] \quad T_{ab} &= 2\sigma_1 l_{(a} m_{b)} \pm l_a l_b + \sigma_2 x_a x_b \\ &\quad + \sigma_3 y_a y_b \\ [31] \quad T_{ab} &= 2\sigma_1 l_{(a} m_{b)} + 2l_{(a} x_{b)} + \sigma_1 x_a x_b \\ &\quad + \sigma_2 y_a y_b \\ [z \bar{z} 11] \quad T_{ab} &= 2\sigma_1 l_{(a} m_{b)} + \sigma_2 (l_a l_b - m_a m_b) \\ &\quad + \sigma_3 x_a x_b + \sigma_4 y_a y_b, \end{aligned} \quad (2)$$

where $\sigma_1, \dots, \sigma_4 \in \mathbb{R}$, having different values for different formulae, and in the Segre class $[z \bar{z} 11]$ $\sigma_2 \neq 0$. In the above "catalog" the first column is the notation commonly used for indicating the Segre class, which is a list $[r_1 r_2 \dots r_n]$ of the dimensions of the Jordan blocks of the corresponding Jordan canonical matrix. The basis vectors $\{l, m, x, y\}$ form a real null tetrad basis such that the only nonvanishing inner products are

$$l^a m_a = x^a x_a = y^a y_a = 1. \quad (3)$$

This basis is constructed from the preferred directions intrinsically defined by the tensor, i.e., from the Jordan basis. The Segre class $[1, 111]$ distinguishes tensors that have diagonal matrix representation and is the unique that admit a timelike eigenvector. The comma in this case is used to separate timelike from spacelike eigenvectors. Regardless of the dimension of a Jordan block, there is only one eigenvector associated to each block, and the eigenvector associated to a block of dimension $r > 1$ is a null vector [4]. Energy-momentum tensors which belong to Segre class $[211]$, for example, have only three linearly independent eigenvectors (one of which is a null vector), and its characteristic polynomial obtained from (1) has three roots: one of multiplicity 2 and the others of multiplicity 1. So, the digits inside the brackets give also the multiplicity of the real eigenvalues, the complex one being taken into account by Segre class $[z \bar{z} 11]$, where the $z \bar{z}$ refers to a pair of complex eigenvalues. Degeneracy amongst eigenvalues in different Jordan blocks will be indicated by enclosing the corresponding digits inside round brackets, as in $[1, (111)]$, which indicate that three out of the four eigenvalues are degenerate. So, to each Segre class there may be several subclasses depending on the degeneracies of the eigenvalues.

A. Segre Classification of Phantom Fields

The energy-momentum tensor for the phantom field has the form [13]

$$T_{ab} = -\phi_a \phi_b + g_{ab} \left[\frac{1}{2} g^{cd} \phi_c \phi_d - V(\phi) \right], \quad (4)$$

where $V(\phi)$ is the phantom potential, $\phi_a \equiv \phi_{;a}$ and the semicolon denotes covariant derivative. Here we assume

that $\phi = \phi(t)$ is a function of time alone evolving in an isotropic and homogeneous space-time, so $g_{ab}\phi^a\phi^b = -\dot{\phi}^2$, that is, ϕ^a is a time-like vector (indeed an eigenvector of T_{ab}). In this case it is always possible to find out two null vectors l_a and m_a such that $l^a m_a = 1$, and ϕ_a can be written as

$$\phi_a = \frac{\dot{\phi}}{\sqrt{2}}(l_a - m_a). \quad (5)$$

Besides, we choose two spacelike vectors x_a and y_a , belonging to the 2-space orthogonal to the 2-space generated by l_a and m_a , so as to form the real null tetrad basis $\{l, m, x, y\}$ defined by (3). In terms of this basis the metric tensor is written as

$$g_{ab} = 2l_{(a}m_{b)} + x_ax_b + y_ay_b \quad (6)$$

where the round brackets indicates symmetrization. Taking into account equations (5) and (6), the canonical form for the energy-momentum tensor (4) is then written as

$$T_{ab} = 2\sigma_1 l_{(a}m_{b)} + \sigma_2 (l_al_b + m_am_b) + \sigma_3 (x_ax_b + y_ay_b), \quad (7)$$

where

$$\sigma_1 = -V(\phi), \quad \sigma_2 = -\frac{1}{2}\dot{\phi}^2, \quad \sigma_3 = -\frac{1}{2}\dot{\phi}^2 - V(\phi), \quad (8)$$

indicating that phantom fields belong to Segre class $[1, 111]$ (see Eq. (2)) for $\sigma_3 = \sigma_4$. In order to find the subclasses we now determine its eigenvalues and corresponding eigenvectors. For the sake of brevity we present our results without going into details of calculations, which can be easily verified from equation (7) and the expressions below. We find that the set of linearly independent eigenvectors and associated eigenvalues of T^a_b is given by

$$\begin{aligned} l^a - m^a &\longrightarrow \left[\frac{1}{2}\dot{\phi}^2 - V(\phi)\right] \\ l^a + m^a &\longrightarrow -\left[\frac{1}{2}\dot{\phi}^2 + V(\phi)\right] \\ x^a &\longrightarrow -\left[\frac{1}{2}\dot{\phi}^2 + V(\phi)\right] \\ y^a &\longrightarrow -\left[\frac{1}{2}\dot{\phi}^2 + V(\phi)\right]. \end{aligned} \quad (9)$$

Note that three out of the four eigenvalues are degenerate, making it apparent that the corresponding Segre subclass is $[1, (111)]$. This is the same Segre type as that for a perfect fluid [16] (we shall comment this latter). We emphasize that this classification is independent of the functional form of the potential $V(\phi)$ as well as the time derivative $\dot{\phi}$ except for $\phi = \text{cte}$. In this latter case the Segre subclass for the phantom field is $[(1, 111)]$ with $-V(\phi)$ a four-time-degenerate eigenvalue. This is the same Segre subclass of energy-momentum tensors of Λ -term type (cosmological constant) [16]. Since a Λ -term can always be incorporated into an energy-momentum tensor of the perfect fluid, we shall banish this subclass from further consideration.

III. CLASSICAL ENERGY CONDITIONS FOR PHANTOM FIELDS

In this Section we investigate possible constraints that the classical energy conditions may impose on the values of the potential $V(\phi)$ as well as on $\dot{\phi}$ for phantom fields. Restrictions imposed by energy conditions on energy-momentum tensors of general relativity theory for matter fields like those represented by Segre classes [211], [31] and $[z\bar{z}11]$ in Eq. (2), have been presented in the literature [1, 17]. As we have found the Segre specializations $[1, (111)]$ and $[(1, 111)]$ for phantom fields, we focus our attention here only on the Segre class $[1, 111]$ of Eq. (2). The most common energy conditions are [17, 18] (see also [19] for a recent discussion on this topic):

- (i) The null energy condition (NEC). NEC states that $T_{ab}n^an^b \geq 0$ for null vectors $n^a \in T_p(M)$ which, for Segre class $[1, 111]$, is equivalent to require that $\sigma_2 - \sigma_1 + \sigma_\alpha \geq 0$ ($\alpha = 3, 4$).
- (ii) The weak energy condition (WEC). WEC states that $T_{ab}t^at^b \geq 0$ for timelike vectors $t^a \in T_p(M)$. This will also imply, by continuity, the NEC. The WEC for Segre class $[1, 111]$ means that $\sigma_2 - \sigma_1 \geq 0$.
- (iii) The strong energy condition (SEC). SEC is the assertion that for any timelike vector $(T_{ab} - T/2g_{ab})t^at^b \geq 0$, where T is the trace of T_{ab} . If T_{ab} belongs to the Segre class $[1, 111]$ then we must have $2\sigma_2 + \sigma_3 + \sigma_4 \geq 0$.
- (iv) The dominant energy condition (DEC). DEC requires that $T_{ab}t^at^b \geq 0$ for timelike vectors $t^a \in T_p(M)$ and the additional requirement that $T_{ab}t^b$ be a non-spacelike vector. By continuity this will also hold for null vectors $n^a \in T_p(M)$. For energy-momentum tensors of Segre class $[1, 111]$ this requires that $\sigma_2 - \sigma_1 \geq 0$, and $\sigma_1 - \sigma_2 \leq \sigma_\alpha \leq \sigma_2 - \sigma_1$ ($\alpha = 3, 4$).

When imposed on the energy-momentum tensor of the phantom field these energy conditions require:

$$\begin{aligned} \text{NEC} &\Rightarrow && -\frac{1}{2}\dot{\phi}^2 \geq 0 \\ \text{WEC} &\Rightarrow && -\frac{1}{2}\dot{\phi}^2 + V(\phi) \geq 0 \\ \text{SEC} &\Rightarrow && \dot{\phi}^2 + V(\phi) \leq 0 \\ \text{DEC} &\Rightarrow && -\dot{\phi}^2 \geq 0 \quad \text{and} \quad V(\phi) \geq 0 \end{aligned}$$

From the above conditions, it should be noticed that although the Segre subclass $[1, (111)]$ for the phantom field is consistent with the Lorentzian signature of the metric tensor g_{ab} , imposition of NEC, as well as DEC, generates at least one contradiction in the sense that the timelike character ($-\dot{\phi}^2 = g_{ab}\phi^a\phi^b$) of the vector ϕ^a is violated. Maintaining the Lorentzian timelike character of ϕ^a WEC can be preserved only for positive potentials since that $V(\phi) \geq \dot{\phi}^2/2$. SEC, on the other hand, cannot be satisfied unless $V(\phi)$ is negative and $V(\phi) \leq -\dot{\phi}^2$. There is however an interval $(-\dot{\phi}^2 < V(\phi) < \dot{\phi}^2/2)$ in




FIG. 1: Energy conditions constraints on phantom fields potential. In interval **(I)** phantom fields do not violate SEC but do violate WEC. In interval **(II)** both (SEC and WEC) are violated while in region **(III)** phantom fields do not violate WEC but do violate SEC.

the values of potential for which neither SEC nor WEC are preserved. Figure 1 shows such intervals. In interval I phantom fields do not violate SEC but do violate WEC. In region II both SEC and WEC are violated while in region III phantom fields do not violate WEC but do violate SEC. It is worth mentioning that evolving quintessence and phantom fields with linear-negative potentials have been recently proposed and studied in light of recent cosmological observations. As shown in Ref. [21], such models provide a good fit for the current supernovae data being, therefore, a potential candidate for dark energy.

At this point, it is important to observe that in our approach the energy conditions refer to quantities like potentials and/or time derivatives of the fields as well as invariantly defined vectors. It is also possible to present the same results in terms of pressures and density associated to the phantom field. As we have noted, the Segre subclass $[1, (111)]$ obtained for the phantom field is the same as that for a perfect fluid. This means that the energy-momentum tensor for an ideal fluid can also be written in the form (7) by identifying $\sigma_1 = (p - \rho)/2$, $\sigma_2 = (p + \rho)/2$ and $\sigma_3 = p$, where p and ρ are the pressure and the density of the fluid.

IV. CONCLUSION

Motivated by a number of observational results, the idea of phantom fields as candidate for the dark energy

has been largely explored in the current literature. In this paper we have examined the algebraic classification of the energy-momentum tensor T_{ab} of such fields for generic potential terms. We have shown that the phantom T_{ab} belongs to Segre subclasses $[1, (111)]$ or $[(1, 111)]$. As is well known, all energy-momentum tensors belonging to these Segre subclasses (e.g., perfect fluid and Λ -term type energy-momentum tensors) couple to Friedmann-Robertson-Walker geometries. We also have found some constraints which are imposed on the values of $V(\phi)$ from the classical energy conditions. Although the SEC is being violated right now, according recent observational data regarding the accelerating universe, we see from our analysis that an evolving potential function $V(\phi)$ can, in principle, be constructed such that this condition is not violated in the past (interval I of FIG. 1), while in that epoch WEC must be violated. Although negative potential for phantom fields have already been proposed, the transitions from interval I to interval II and to III demand some studies. Our approach makes also clear that imposition of NEC, as well as DEC, on phantom fields generates a strong contradiction about the timelike character of the vector ϕ^a composing the energy-momentum tensor.

As discussed earlier, the Segre classification technique can also be applied to the geometric part of Einstein's equations. In this case, the results could be used to impose the energy conditions directly, via Segre classes, on the coefficients of a given metric such as, for example, the FRW metric. From such an analysis, the production of future singularities in an expanding universe without violating the classical energy conditions could be studied (see, e.g. [22]). This possibility will be explored in a forthcoming communication.

Acknowledgments

The authors are very grateful to M. J. Rebouças for valuable discussions. JSA thanks the Department of Physics of the Universidade Federal do Rio Grande do Norte for their hospitality during the course of this work, and also the financial support by CNPq (62.0053/01-1-PADCT III/Milenio). JS thanks the support of PRONEX/CNPq/FAPERN.

-
- [1] G. S. Hall, *Diff. Geom.* **12**, 53 (1983). This reference contains an extensive bibliography on the algebraic classification of a second order tensor on 4-dimensional Lorentzian space-times manifolds.
[2] R. V. Churchill, *Trans. Amer. Math. Soc.* **34**, 784 (1932);

- W. J. Cormack and G. S. Hall, *J. Phys. A* **12**, 55 (1979).
[3] G. S. Hall and D. A. Negm, *Int. J. Theor. Phys.* **25**, 405 (1986); J. Santos, M. J. Rebouças and A. F. F. Teixeira, *J. Math. Phys.* **34**, 186 (1993).
[4] J. Santos, M.J. Rebouças and A.F.F. Teixeira, *J. Math.*

- Phys.* **36**, 3074 (1995); J. Santos, M.J. Rebouças and A.F.F. Teixeira, *Gen. Rel. Grav.* **27**, 989 (1995).
- [5] M. J. Rebouças and J. Santos, *Mod. Phys. Lett. A* **18**, 2807 (2003).
- [6] A. Karlhede, *Gen. Rel. Grav.* **12**, 693 (1980); M. A. H. MacCallum, “Computer-aided Classification of Exact Solutions in General Relativity”, in *General Relativity and Gravitational Physics (9th Italian Conference)*, edited by R. Cianci, R. de Ritis, M. Francaviglia, G. Marmo, C. Rubano and P. Scudellaro. World Scientific Publishing Co., Singapore (1991).
- [7] D. N. Spergel et al., *Astrophys. J. Suppl.* **148**, 175 (2003)
- [8] A. G. Riess et al., *Astrophys. J.* **607** 665 (2004)
- [9] V. Sahni and A. Starobinsky, *Int. J. Mod. Phys. D* **9**, 373 (2000); J. E. Peebles and B. Ratra *Rev. Mod. Phys.* **75**, 559 (2003); T. Padmanabhan, *Phys. Rept.* **380**, 235 (2003); J. A. S. Lima, *Braz. J. Phys.* **34**, 194 (2004).
- [10] B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37**, 3406 (1988); R. R. Caldwell, R. Dave, and P. J. Steinhardt, *Phys. Rev. Lett.* **80**, 1582 (1998); P. G. Ferreira and M. Joyce, *Phys. Rev. D* **58**, 023503 (1998)
- [11] R. R. Caldwell, *Phys. Lett. B* **545**, 23 (2002); B. McInnes, *astro-ph/0210321*; V. Faraoni, *Int. J. Mod. Phys. D* **11**, 471 (2002); S. M. Carroll, M. Hoffman and M. Trodden, *Phys. Rev. D* **68**, 023509 (2003); S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **562**, 147 (2003); J. S. Alcaniz, *Phys. Rev. D* **69**, 083521 (2004); J. A. S. Lima and J. S. Alcaniz, *Phys. Lett. B* **600**, 191 (2004); J. M. Aguirregabiria, L. P. Chimento and R. Lazkoz, *Phys. Rev. D* **70**, 023509 (2004)
- [12] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, *Phys. Rev. Lett.*, **91**, 071301 (2003); P. F. Gonzalez-Diaz, *Phys. Rev. D* **68**, 021303 (2003); S. Nesseris and L. Perivolaropoulos, *Phys. Rev. D* **70**, 123529 (2004); P. F. Gonzalez-Diaz and J. A. Jimenez-Madrid, *Phys. Lett. B* **596**, 16 (2004).
- [13] P. Singh, M. Sami, and N. Dadhich, *Phys. Rev. D* **68**, 023522 (2003).
- [14] T. Chiba, T. Okabe and M. Yamaguchi, *Phys. Rev. D* **62**, 023511 (2000)
- [15] V. Sahni and Y. Shtanov, *Int. J. Mod. Phys. D* **11**, 1515 (2002); *J. Cosm. Astrop. Phys.* **0311**, 014 (2003)
- [16] D. Kramer, H. Stephani, E. Herlt and M. MacCallum, *Exact Solutions of Einstein's Field Equations*, Cambridge U. P., Cambridge (1980), pg. 67.
- [17] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Spacetime*, (Cambridge University Press, England, 1973).
- [18] M. Visser, *Lorentzian Wormholes*, (AIP Press, New York, 1996).
- [19] M. Visser, *Science* **276**, 88 (1997); M. Visser and C. Barceló, “Energy conditions and their cosmological implications”, *Cosmo99 Proceedings: Trieste, Sept/Oct 1999*. World Scientific; gr-qc/0001099; C. Barceló and M. Visser, *Int. J. Mod. Phys. D* **11**, 1553 (2002).
- [20] M. J. Rebouças, J. Santos and A. F. F. Teixeira, *Braz. J. Phys.* **34**, 535 (2004).
- [21] L. Perivolaropoulos, Preprint *astro-ph/0412308*.
- [22] J.D. Barrow, *Class. Quantum Grav.* **21**, L79 (2004); J.D. Barrow, *Class. Quantum Grav.* **21**, 5619 (2004), J.D. Barrow and C.G. Tsagas, Preprint *gr-qc/0411045*